

**Page 34 Line 9.** **After** summer program. **add** Exercise 1.6.3 and 1.6.4 are stated as “either/or”. It is possible that *both* alternatives occur: there is some number with more than one expansion, while there is another number with no expansion. For example, modify the Eisenstein number system (Exercise 1.6.10) by using base  $b = -2$ , but using only 3 of the digits, say  $D = \{0, 1, \omega\}$ . (Thanks to Peter Hinow for this example.)

**Page 43 Line 6.** **Replace** open set  $U$  **by** open set  $T$

**Page 43 Line 7.** **Replace**  $U = \bigcup_{A \in \mathcal{A}} A$  **by**  $T = \bigcup_{U \in \mathcal{A}} U$

**Page 84 Line 7.** **After** 3.1.1. **add** Let  $W$  and  $F$  be as on p. 83.

**Page 96 Line 5.** **Replace** no cover of  $\mathbb{R}^2$  by bounded open sets with order 1.  
**by** no cover with order 1 of  $\mathbb{R}^2$  by open sets with bounded diameter.

**Page 97 Line 9.** **Replace**  $F \cup \bigcup_{i=1}^{n+1} B_i$  **by**  $F_1 \cup \bigcup_{i=1}^{n+1} B_i$

**Page 97 Line -14.** **Replace** Theorem 3.4.2 **by** Theorem 3.4.3

**Page 100 Line 12.** **Replace Exercise by**

By Theorems (3.4.4) and (3.4.5), we may conclude that  $\text{Cov } \mathbb{R} = 1$  since  $\text{ind } \mathbb{R} = 1$ . Instead, compute  $\text{Cov } \mathbb{R}$  directly from the definition of covering dimension.

**Page 114 Line 10.** **After** Mauldin–Williams graph

**add** For technical reasons we assume each vertex has at least one edge leaving it.

**Page 115 Line -8.** **Replace** complete metric **by** nonempty complete metric

**Page 129 Line -18.** **Replace two paragraphs beginning** Now we take **by**

Now we take the case  $\mathcal{L}(A) = \infty$ . All of the sets  $A \cap [-n, n]$  are measurable, so there exist open sets  $U_n \supseteq A \cap [-n, n]$  and compact sets  $F_n \subseteq A \cap [-n, n]$  with  $\mathcal{L}(U_n \setminus F_n) < \varepsilon/2^{n+2}$ . Define  $U'_n = U_n \cap ((-\infty, -n+1 + \varepsilon/2^{n+2}) \cup (n-1 - \varepsilon/2^{n+2}, \infty))$  and  $F'_n = F_n \cap ([-n, -n+1] \cup [n-1, n])$ , so that  $U'_n$  is open,  $F'_n$  is compact,  $U'_n \supseteq A \cap ([-n, -n+1] \cup [n-1, n]) \supseteq F'_n$  and  $\mathcal{L}(U'_n \setminus F'_n) < 3\varepsilon/2^{n+2} < \varepsilon/2^n$ . Now  $U = \bigcup U'_n$  is open, and  $F = \bigcup F'_n$  is closed (Exercise 2.1.42). We have  $U \supseteq A \supseteq F$ , and  $U \setminus F \subseteq \bigcup_{n \in \mathbb{N}} (U'_n \setminus F'_n)$ , so that  $\mathcal{L}(U \setminus F) \leq \sum \mathcal{L}(U'_n \setminus F'_n) < \varepsilon$ .

Conversely, suppose sets  $U$  and  $F$  exist. By Theorem (5.1.12) they are measurable. First assume  $\overline{\mathcal{L}}(A) < \infty$ . Then  $\mathcal{L}(F) < \infty$ , and  $\mathcal{L}(U) \leq \mathcal{L}(U \setminus F) + \mathcal{L}(F) < \varepsilon + \mathcal{L}(F) < \infty$ . Now  $\overline{\mathcal{L}}(A) \leq \overline{\mathcal{L}}(U) = \mathcal{L}(U) < \mathcal{L}(F) + \varepsilon = \underline{\mathcal{L}}(F) + \varepsilon \leq \underline{\mathcal{L}}(A) + \varepsilon$ . This is true for any  $\varepsilon > 0$ , so  $\overline{\mathcal{L}}(A) = \underline{\mathcal{L}}(A)$ , so  $A$  is measurable.

**Page 134 Line 10.** **Add** (Recall that  $\inf \emptyset = \infty$ , so if there is no countable cover at all of  $B$  by sets of  $\mathcal{A}$ , then  $\overline{\mathcal{M}}(B) = \infty$ .)

**Page 154 Line 16.** **Replace**  $h(a)$  **by**  $h(f(a))$  **and replace**  $h(b)$  **by**  $h(f(b))$

**Page 160 Line 18.** **After** system **add** on a complete metric space

**Page 161 Line 16.** **After** (6.3.11) **add** Let  $f_i$  and  $K$  be as in Theorem (4.1.3)

**Page 190 Line -10.** **Replace** let **by** Let

**Page 208 Line 7.** **Replace** Let **by** For  $\sigma \in E^{(\omega)}$ , let

**Page 213 Line 20.** **Replace**  $-\log a / \log 5$  where  $a \approx 0.3992$  is a solution of  $a^4 - a^3 - a^2 + 3a - 1 = 0$ . The dimension is approximately 0.57058;

**by**  $\log((\sqrt{5} + 3)/2) / \log 5 \approx 0.59799$ ;

**Changes: from second printing to third printing** February 7, 1995  
**Measure, Topology, and Fractal Geometry** by Gerald A. Edgar

**Page 68 Line 10.** Replace The points of  $F$  by The points of  $F_1$

**Page 83 Line 8.** After  $\{U_1, U_2, \dots\}$ .

**add** (If  $\mathcal{B}$  is finite, repeat the basic sets over and over.)

**Page 109 Line 4.** Replace complex number by complex number,  $|b| > 1$

**Page 145, after line 3.** Add

An example of a set in the line not measurable for Lebesgue measure may be found in many texts. For example: [6, pp. 36–37], [9, Theorem 1.4.7], [29, (10.28)], or [48, Chapter 3, Section 4].

**Page 187 Line -14.** Replace Theorem 6.5.2 by Theorem 6.2.5

**Changes: from first printing to second printing** March 28, 1992  
**Measure, Topology, and Fractal Geometry** by Gerald A. Edgar

**Page ix Line 14.** Delete and  
**Page ix Line 15.** Add and Kenneth Dreyhaupt  
**Page 4 Line 23.** Replace 1.6.7 by 1.6.2  
**Page 17 Line 8.** Replace 24 by 2.4  
**Page 29 last display.**

Replace  $(2\pi/3)$  by  $\frac{2\pi}{3}$

**Page 29–30.** Replace  $a$  and  $b$  by  $u$  and  $v$   
**Page 31 Figure 1.6.8.** Missing label  $L_4$ .  
**Page 32 Line -9.** Replace is has come by it has come  
**Page 33 Line 2.**

Replace called by Mandelbrot “Sierpiński’s carpet”  
 by called Sierpiński’s carpet (*dywan Sierpinskiego*)

**Page 49 Line -10.**

Replace  $\{y \in \mathbb{R}^d : \rho(x, y) = r\}$  by  $\{y \in S : \rho(x, y) = r\}$

**Page 57 Line 4.** Replace  $\rightarrow z_1$  by  $= z_1$

**Page 57 Line 7.** Replace  $\rightarrow z_2$  by  $= z_2$

**Page 57 Line 9.** Replace  $\rightarrow z_j$  by  $= z_j$

**Page 65 Line 5.** Replace 2.3.15 by 2.3.16

**Page 66 Line 19.** Replace first paragraph of proof by

PROOF. First, clearly  $D(A, B) \geq 0$  and  $D(A, B) = D(B, A)$ . Since  $A$  and  $B$  are compact, they are bounded, so  $D(A, B) < \infty$ .

**Page 66 Line -3.** Add and let  $A$  be a nonempty compact subset of  $S$

**Page 67–68. Replace (2.4.4) by**

**(2.4.4) THEOREM.** *Suppose  $S$  is a complete metric space. Then the space  $\mathcal{K}(S)$  is complete.*

PROOF. Suppose  $(A_n)$  is a Cauchy sequence in  $\mathcal{K}(S)$ . I must show that  $A_n$  converges. Let

$$A = \{x : \text{there is a sequence } (x_k) \text{ with } x_k \in A_k \text{ and } x_k \rightarrow x\}.$$

I must show that  $D(A_n, A) \rightarrow 0$  and  $A$  is nonempty and compact.

Let  $\varepsilon > 0$  be given. Then there is  $N \in \mathbb{N}$  so that  $n, m \geq N$  implies  $D(A_n, A_m) < \varepsilon/2$ . Let  $n \geq N$ . I claim that  $D(A_n, A) \leq \varepsilon$ .

If  $x \in A$ , then there is a sequence  $(x_k)$  with  $x_k \in A_k$  and  $x_k \rightarrow x$ . So, for large enough  $k$ , we have  $\rho(x_k, x) < \varepsilon/2$ . Thus, if  $k \geq N$ , then (since  $D(A_k, A_n) < \varepsilon/2$ ) there is  $y \in A_n$  with  $\rho(x_k, y) < \varepsilon/2$ , and we have  $\rho(y, x) \leq \rho(y, x_k) + \rho(x_k, x) < \varepsilon$ . This shows that  $A \subseteq N_\varepsilon(A_n)$ .

Now suppose  $y \in A_n$ . Choose integers  $k_1 < k_2 < \dots$  so that  $k_1 = n$  and  $D(A_{k_j}, A_m) < 2^{-j}\varepsilon$  for all  $m \geq k_j$ . Then define a sequence  $(y_k)$  with  $y_k \in A_k$  as follows: For  $k < n$ , choose  $y_k \in A_k$  arbitrarily. Choose  $y_n = y$ . If  $y_{k_j}$  has been chosen, and  $k_j < k \leq k_{j+1}$ , choose  $y_k \in A_k$  with  $\rho(y_{k_j}, y_k) < 2^{-j}\varepsilon$ . Then  $y_k$  is a Cauchy sequence, so it converges. Let  $x$  be its limit. So  $x \in A$ . We have  $\rho(y, x) = \lim_k \rho(y, y_k) < \varepsilon$ . So  $y \in N_\varepsilon(A)$ . This shows that  $A_n \subseteq N_\varepsilon(A)$ . Note that, taking  $\varepsilon = 1$  in this argument, I have also proved that  $A \neq \emptyset$ .

So we have  $D(A, A_n) \leq \varepsilon$ . This concludes the proof that  $(A_n)$  converges to  $A$ .

Next I show that  $A$  is “totally bounded”: that is, for every  $\varepsilon > 0$ , there is a finite  $\varepsilon$ -net in  $A$ . Choose  $n$  so that  $D(A_n, A) < \varepsilon/3$ . By (2.2.5), there is a finite  $(\varepsilon/3)$ -net for  $A_n$ , say  $\{y_1, y_2, \dots, y_m\}$ . Now for each  $y_i$ , there is  $x_i \in A$  with  $\rho(x_i, y_i) < \varepsilon/3$ . The finite set  $\{x_1, x_2, \dots, x_m\}$  is an  $\varepsilon$ -net for  $A$ .

Now I will show that  $A$  is a closed subset of  $S$ . Let  $x$  belong to the closure  $\bar{A}$  of  $A$ . Then there exists a sequence  $(y_n)$  in  $A$  with  $\rho(x, y_n) < 2^{-n}$ . For each  $n$  there is a point  $z_n \in A_n$  with  $\rho(z_n, y_n) < D(A_n, A) + 2^{-n}$ . Now

$$\rho(z_n, x) \leq \rho(z_n, y_n) + \rho(y_n, x) < D(A_n, A) + 2^{-n} + 2^{-n}.$$

This converges to 0, so  $z_n \rightarrow x$ . Thus  $x \in A$ . This shows that  $A$  is closed.

Finally, to show that  $A$  is compact, I will show that it is countably compact. Let  $F$  be an infinite subset of  $A$ . There is a finite  $(1/2)$ -net  $B$  for  $A$ , so each element of  $F$  is within distance  $1/2$  of some element of  $B$ . Now  $F$  is infinite and  $B$  is finite, so there is an element of  $B$  within distance  $1/2$  of infinitely many elements of  $F$ . Let  $F_1 \subseteq F$  be that infinite subset. The points of  $F_1$  are all within distance  $1$  of each other; that is,  $\text{diam } F_1 \leq 1$ . In the same way, there is an infinite set  $F_2 \subseteq F_1$  with  $\text{diam } F_2 \leq 1/2$ ; and so on. There are infinite sets  $F_j$  with  $\text{diam } F_j \leq 2^{-j}$  and  $F_{j+1} \subseteq F_j$  for all  $j$ . Now if  $x_j$  is chosen from  $F_j$ , we have  $\rho(x_j, x_k) \leq 2^{-j}$  if  $j < k$ , so  $(x_j)$  is a Cauchy sequence. Since  $S$  is complete,  $(x_j)$  converges, say  $x_j \rightarrow x$ . Since  $A$  is closed,  $x \in A$ . But then  $x$  is a cluster point of the set  $F$ . Therefore  $A$  is compact. ☺

**Page 74 headline.** Replace 2.5.12 by 2.5.12

**Page 76 Line -6.**

**Replace** Then apply 2.2.18

**by** But  $A$  is closed, so any point with distance 0 from  $A$  belongs to  $A$ .

**Page 86 Line -15. After** empty set. **add**

Note that a space  $S$  has  $\text{ind } S \leq k$  if and only if a point  $\{x\}$  and a closed set  $B$  not containing  $x$  can be separated by a set  $L$  with  $\text{ind } L \leq k - 1$ . Indeed, there is a basic open set  $U$  with  $x \in U \subseteq S \setminus B$  and  $L = \partial U$  separates  $\{x\}$  and  $B$ .

**Page 92 Line -1. Replace**  $(ay + bx)$  **by**  $(ay - bx)$

**Page 93 Line -10. Delete comma**

**Page 94 Line -12. Replace**  $g_1(x) \geq x_2 \geq -1$  **by**  $g_1(x) \geq x_1 \geq -1$

**Page 98 Line -3. Replace** finite paths **by** of finite paths

**Page 102 Line -4. Replace**  $(i = 1, 2, \dots, m)$  **by**  $(i = 1, 2, \dots, n)$

**Page 103 Line -16. Replace** Exercise 3.1.7 **by** Exercise 3.1.8

**Page 108 Line -7. Replace** a cluster of **by** a cluster point of

**Page 109 Line 5. Replace**  $\dots d_k$  **by**  $\dots, d_k$

**Page 113 Line 7. After** similarities. **add** (See Plate 8.)

**Page 113 Line -1. After** made. **add**

The transformations map the large square and rectangle shown at the top into the same square and rectangle shown at the bottom.

**Page 114 Line 9. Replace**  $r: V \rightarrow (0, \infty)$  **by**  $r: E \rightarrow (0, \infty)$

**Page 117 Line 5. Replace paragraph by**

Under the new metrics, what happens to the maps  $f_e$ ? If  $e \in E_{uv}$ , then

$$\begin{aligned} \rho'_u(f_e(x), f_e(y)) &= a_u \rho_u(f_e(x), f_e(y)) \\ &= a_u r(e) \rho_v(x, y) \\ &= \frac{a_u r(e)}{a_v} \rho'_v(x, y). \end{aligned}$$

Thus, with the new metrics, the maps  $f_e$  realize a Mauldin-Williams graph  $(V, E, i, t, r')$ , where

$$r'(e) = \frac{a_u}{a_v} r(e) \quad \text{for } e \in E_{uv}.$$

The Mauldin-Williams graph  $(V, E, i, t, r')$  is called a **rescaling** of the graph  $(V, E, i, t, r)$ . A Mauldin-Williams graph  $(V, E, i, t, r)$  will be called **contracting** iff it is a rescaling of a strictly contracting graph.

**Page 126 Line 5. Replace** 1.5.2 **by** 5.1.2

**Page 127 Line 11.**

**Replace** and let  $A \subseteq \bigcup_{j \in \mathbb{N}} [a_j, b_j)$  **by** and let  $A \cup B \subseteq \bigcup_{j \in \mathbb{N}} [a_j, b_j)$

**Page 129. Replace statement of Theorem (5.1.12) by**

*Compact subsets, closed subsets, and open subsets of  $\mathbb{R}$  are Lebesgue measurable.*

**Page 129 Line 4. Replace the first paragraph of the proof by**

PROOF. Let  $K \subseteq \mathbb{R}$  be compact. It is bounded, so  $K \subseteq [-n, n]$  for some  $n$ , and therefore  $\overline{\mathcal{L}}(K) < \infty$ . The compact set  $K$  is a subset of  $K$ , so  $\underline{\mathcal{L}}(K) \geq \overline{\mathcal{L}}(K)$ .

Let  $F \subseteq \mathbb{R}$  be a closed set. Then for each  $n \in \mathbb{N}$ , the intersection  $F \cap [-n, n]$  is compact, and therefore measurable. Thus  $F$  is measurable.

**Page 131 Line 15. Replace  $\overline{\mathcal{L}}(f[U] \setminus f[F]) < r\varepsilon$  by  $\overline{\mathcal{L}}(f[U] \setminus f[F]) = \overline{\mathcal{L}}(f[U \setminus F]) < r\varepsilon$**

**Page 135 Line 12.**

**(wrong font) Replace  $\overline{\mathcal{M}}$ -measurable by  $\overline{\mathcal{M}}$ -measurable**

**Page 135 Line -3. Replace period by comma**

**Page 140 Line 13. Replace let  $(f_1, f_2, \dots, f_n)$  is a by let  $(f_1, f_2, \dots, f_n)$  be a**

**Page 154 Line 2. Replace  $\varepsilon/2$  by  $\varepsilon$**

**Page 154 Line 10.**

**Replace  $0 \leq y_1 < z_1 \leq y_2 < z_2 \leq \dots \leq y_n < z_n \leq 1$**

**by  $0 \leq y_1 \leq z_1 \leq y_2 \leq z_2 \leq \dots \leq y_n \leq z_n \leq 1$**

**Page 154 Line 19.**

**Replace  $\rho(u, v)$ , so we have by  $\rho(u, v)$ , so  $h$  has bounded increase, and we have**

**Page 154 Line -6. Replace then the sets  $f[[x_{i-1}, x_i]]$  are measurable and disjoint by then the set  $f[[x_{i-1}, x_i]] = f[[x_{i-1}, x_i]] \setminus \{f(x_i)\}$  is the difference of two compact sets, hence measurable. The sets  $f[[x_{i-1}, x_i]]$  are disjoint**

**Page 155 Line 3. On Hausdorff dimension vs. topological dimension add a footnote:**

*\*Optional material. (It depends on Section 3.4.)*

**Page 156 Line 20. After system. add Of course,  $K$  is a measurable set, since it is compact.**

**Page 163 Line 9. Replace *conditon* by *condition***

**Page 180. Refer to Tricot as well as Taylor.**

**Page 186 Line 4. Replace A. N. Besicovitch by A. S. Besicovitch**

**Page 191 Line -2.**

**Replace inside of the fudgeflake by filled-in fudgeflake**

**Page 192 Line 12. Replace Exercise 4.3.12 by Figure 4.3.12**

**Page 224 Line -18. Replace Kurt by Curt**

**Page 228. Replace Maltese cross xii by Maltese cross xiii**

**Page 228.**

**Replace number systems 28, 34, 59, 83, 34, 109, 131, 165, 178, 204**

**by number systems 28, 34, 59, 83, 109, 131, 165, 177, 204**

**Page 228. Space between overlap and packing dimension**