

(1.) Let $A = \begin{pmatrix} 3 & 6 & 3 & 3 & 6 & 1 \\ 5 & 10 & 4 & 2 & -3 & 6 \\ 7 & 14 & 2 & 0 & -11 & 7 \\ 1 & 2 & 0 & 0 & -1 & 2 \end{pmatrix}$ and find:

- (a.) A basis for the row space of A .
- (b.) A basis for the column space of A .
- (c.) A basis for the nullspace of A . DO NOT USE MATLAB.
- (d.) The rank of A .

(2.) Suppose that $\mathbf{v}_1 = \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix}$, $\mathbf{v}_2 = \begin{pmatrix} 0 \\ -1 \\ 3 \end{pmatrix}$, $\mathbf{v}_3 = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$ and $\mathbf{v}_4 = \begin{pmatrix} 2 \\ 2 \\ 8 \end{pmatrix}$. Find a basis for $\text{Span}(\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4)$.

Note: there is more than one answer.

(3.) Determine if S is a subspace of V where:

- (a.) $V = \mathbb{R}^{2 \times 2}$ and S is the set of 2×2 matrices A with $\det(A)=0$.
- (b.) $V = \mathbb{R}^{2 \times 2}$ and S is the set of 2×2 upper triangular matrices.
- (c.) $V = \mathbb{R}^2$ and $S = \{(x_1, x_2)^T \mid |x_1| = |x_2|\}$.
- (d.) $V = \mathbb{P}_5$ and S is the set of all polynomials $p(x)$ in V such that $p(1) = 0$.
- (e.) $V = C[-1, 1]$ and S is the set of odd functions in V .

(4.) Determine if $1, e^x$ and $\cos x$ are linearly independent in $C[0, 1]$.

(5.) Find a basis for the subspace S of V where:

- (a.) $V = \mathbb{R}^4$ and $S = \{(a - b + c, a + c, a + 2b - c, b - 3c)^T \mid a, b, c \text{ are real numbers}\}$.
- (b.) $V = C[0, 1]$ and $S = \text{Span}(1, \sin 2x, \sin x \cos x)$.
- (c.) $V = \mathbb{P}_4$ and S is the set of all polynomials $p(x)$ in V with $p(0) = 0$ and $p(1) = 0$.

(6.) Let $A = \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}$ and let $S = \{B \in \mathbb{R}^{2 \times 2} \mid AB = BA\}$.

- (a.) Show that S is a subspace of $\mathbb{R}^{2 \times 2}$.
- (b.) Find a basis for S .

(7.) True or False?

- (a.) A linearly independent set can not contain $\mathbf{0}$.
- (b.) A subspace of a vector space must contain $\mathbf{0}$.
- (c.) If A is an $m \times n$ matrix, then $\dim(\text{Col}(A)) + \dim(\text{N}(A)) = m$.
- (d.) If A is an $m \times n$ matrix, then $A\mathbf{x} = \mathbf{b}$ is consistent if and only if \mathbf{b} is in the column space of A .
- (e.) If A is a singular $n \times n$ matrix, then the columns of A form a basis for \mathbb{R}^n .
- (f.) A spanning set can never be linearly independent.

(8.) Prove the following:

(a.) Let S be the subset of $\mathbb{R}^{n \times n}$ consisting of matrices A such that $A^T = A$. Show that S is a subspace of $\mathbb{R}^{n \times n}$. (The matrices in S are called symmetric matrices.)

(b.) If $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ are linearly independent vectors in a vector space V , then $\{\mathbf{v}_2, \dots, \mathbf{v}_n\}$ do not span V .

(c.) If $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ are linearly independent vectors in a vector space V which do not span V , then there is a vector \mathbf{v}_0 in V such that $\{\mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_n\}$ is also linearly independent.