

Review 1

(1.) Consider the following system of linear equations:

$$\begin{aligned} 3x_1 - 2x_2 + 4x_3 + 3x_4 - x_5 &= 17 \\ 2x_1 + 4x_3 + x_4 + 2x_5 &= 8 \\ -2x_1 + 2x_2 - 2x_3 - x_4 + 2x_5 &= -10 \\ -2x_1 - 2x_2 - 6x_3 + 2x_4 - 6x_5 &= 0 \end{aligned}$$

- (a.) Find the augmented matrix of this system.
- (b.) Find the reduced row echelon form of the matrix in (a.).
- (c.) Is this system consistent? If so, find all solutions.

(2.) Given the augmented matrix $A = \left(\begin{array}{ccc|c} 1 & -1 & 3 & 1 \\ 3 & 2 & -2 & -2 \\ -2 & 7 & \alpha & \beta \end{array} \right)$, find all values of α and β such that the corresponding linear system has:

- (a.) No solutions.
- (b.) A unique solution.
- (c.) Infinitely many solutions.

(3.) Determine if S is a subspace of V where:

- (a.) $V = \mathbb{R}^{2 \times 2}$ and S is the set of 2×2 matrices A with $\det(A) = 0$.
- (b.) $V = \mathbb{R}^{2 \times 2}$ and S is the set of 2×2 upper triangular matrices.
- (c.) $V = \mathbb{R}^2$ and $S = \{(x_1, x_2)^T \mid |x_1| = |x_2|\}$.
- (d.) $V = \mathbb{P}_5$ and S is the set of all polynomials $p(x)$ in V such that $p(1) = 0$.
- (e.) $V = C[-1, 1]$ and S is the set of odd functions in V .

(4.) Suppose $A = \begin{pmatrix} 2 & 1 & 3 \\ 0 & 0 & 7 \\ 4 & 0 & 11 \end{pmatrix}$ and $B = \begin{pmatrix} 2 & 1 & 3 \\ 0 & -2 & 5 \\ 0 & 0 & 1 \end{pmatrix}$

- (a.) Find three elementary matrices \mathcal{E}_1 , \mathcal{E}_2 , and \mathcal{E}_3 such that $\mathcal{E}_3\mathcal{E}_2\mathcal{E}_1A = B$.
- (b.) Compute $\det(A)$ using the results of part (a.). DO NOT USE MATLAB.

(5.) Find an LU -factorization for $A = \begin{pmatrix} 1 & 3 & 5 \\ 2 & 4 & 7 \\ -2 & 4 & 1 \end{pmatrix}$. DO NOT USE MATLAB

(6.) True or False. You do not need to explain your answer.

- (a.) If A and B are $n \times n$ matrices, then $\det(A + B) = \det(A) + \det(B)$.
- (b.) If A and B are $n \times n$ matrices, then $AB = BA$.
- (c.) Suppose $A\mathbf{x} = \mathbf{b}_1$ has a unique solution, then it is possible for $A\mathbf{x} = \mathbf{b}_2$ to have more than one solution.
- (d.) An underdetermined system of linear equations is always consistent.
- (e.) Every $n \times n$ elementary matrix is invertible.
- (f.) If A is an $n \times n$ invertible matrix, then A can be row reduced to I_n .
- (g.) If A is an $n \times n$ matrix and α is a scalar, then $\det(\alpha A) = \alpha \det(A)$.

(7.) Prove the following:

- (a.) If A is an invertible, then $\det(A^{-1}) = \det(A)^{-1}$.
- (b.) If A is an invertible, then $A\mathbf{x} = \mathbf{b}$ has a unique solution for each \mathbf{b} .
- (c.) If A is an invertible, then $A = \mathcal{E}_1\mathcal{E}_2 \cdots \mathcal{E}_k$ where $\mathcal{E}_1, \mathcal{E}_2, \dots, \mathcal{E}_k$ are some elementary matrices.
- (d.) If $A\mathbf{x}_1 = \mathbf{b}$ and $A\mathbf{x}_2 = \mathbf{b}$, then $(\mathbf{x}_1 - \mathbf{x}_2)$ is a solution to $A\mathbf{x} = \mathbf{0}$.