

# Week 9 Lectures, Math 716, Tanveer

## 1 Green's function as a distribution

### 1.1 Laplace Operator

For the Poisson-Problem with homogeneous boundary condition:

$$\Delta u = f \text{ for } \mathbf{x} \in \Omega, \quad u = 0 \text{ on } \partial\Omega \quad (1)$$

we know that

$$u(\mathbf{x}_0) = \int_{\Omega} G(\mathbf{x}, \mathbf{x}_0) f(\mathbf{x}) d\mathbf{x} \quad (2)$$

On the otherhand, if  $u$  is a test function with support inside  $\Omega$ , we have from using corollary 4 of week 8 notes that

$$u(\mathbf{x}_0) = \int_{\Omega} G(\mathbf{x}, \mathbf{x}_0) \Delta u(\mathbf{x}) d\mathbf{x} = (u, \Delta G(\cdot, \mathbf{x}_0)) \quad (3)$$

Therefore, in the sense of distribution,

$$\Delta G(\mathbf{x}, \mathbf{x}_0) = \delta(\mathbf{x} - \mathbf{x}_0) \quad (4)$$

Therefore, we view solution (2) as a principle of linear superposition. In the physical context ( $n = 3$ ), it means that the potential caused by charge density  $f$  in a domain  $\Omega$  with boundary at zero potential is given by a linear superposition of point charge potentials satisfying the same boundary conditions, with a weighting proportional to the infinitesimal charge  $f(\mathbf{x})d\mathbf{x}$  present in a volume element  $d\mathbf{x}$  at  $\mathbf{x}$ .

Further, note that  $G(\mathbf{x}, \mathbf{x}_0) = G_0(|\mathbf{x} - \mathbf{x}_0|) + H(\mathbf{x}, \mathbf{x}_0)$ , where  $H$  is harmonic in  $\mathbf{x}$ . It follows that

$$\Delta G_0(|\mathbf{x} - \mathbf{x}_0|) = \delta(\mathbf{x} - \mathbf{x}_0) \quad (5)$$

### 1.2 Heat Equation

Recall from last class that the source function:

$$\mathcal{S} = \left( \frac{1}{4\pi\kappa t} \right)^{n/2} \exp \left[ -\frac{|\mathbf{x}|^2}{4\kappa t} \right] \quad (6)$$

satisfies

$$\mathcal{S}_t = \kappa \Delta \mathcal{S} \text{ for } \mathbf{x} \in \mathbb{R}^n, t > 0, \text{ with } \mathcal{S}(\mathbf{x}, 0^+) = \delta(\mathbf{x}) \quad (7)$$

It can be shown (exercise) that

$$R(\mathbf{x}, t) = \mathcal{S}(\mathbf{x} - \mathbf{x}_0, t - t_0) \text{ for } t > t_0 \text{ and } R(\mathbf{x}, t) = 0 \text{ for } t < t_0 \quad (8)$$

satisfies

$$R_t - \kappa \Delta R = \delta(\mathbf{x} - \mathbf{x}_0) \delta(t - t_0) \quad (9)$$

## 2 Wave equation

### 2.1 Solution in higher dimension through Spherical Means

Assume  $u$  is a classical solution to the initial value problem for  $n$ -dimensional wave equation for  $n \geq 2$ :

$$u_{tt} - \Delta u = 0 \text{ for } \mathbf{x} \in \mathbb{R}^n \text{ for } t > 0 \text{ with } u(\mathbf{x}, 0) = \phi(\mathbf{x}), \quad u_t(\mathbf{x}, 0) = \psi(\mathbf{x}) \quad (10)$$

where  $\phi \in \mathbf{C}^2$  and  $\psi \in \mathbf{C}^1$ . For  $t > 0, r > 0$ , we define  $U(\mathbf{x}; r, t)$  to be the spherical average over the surface of an  $n$ -dimensional ball  $B(\mathbf{x}, r)$  of radius  $r$ , centered at  $\mathbf{x}$ , and denoted by

$$U(\mathbf{x}; r, t) = \frac{1}{A_r} \int_{\partial B(\mathbf{x}; r)} u(\mathbf{y}, t) d\mathbf{y} \equiv \int_{\partial B(\mathbf{x}; r)} u(\mathbf{y}, t) d\mathbf{y} \quad (11)$$

where  $A_r$  is the surface area of an  $n$  dimensional ball of radius  $r$ . Note  $A_r = n\alpha(n)r^{n-1}$ , where volume of the  $n$ -dimensional sphere is  $\alpha(n)r^n$ . It is to be noted that

$$\lim_{r \rightarrow 0^+} U(\mathbf{x}; r, t) = u(\mathbf{x}, t)$$

from continuity of  $\mathbf{u}$ . We can similarly define

$$G(\mathbf{x}; r) = \int_{\partial B(\mathbf{x}; r)} \phi(\mathbf{y}, t) d\mathbf{y} \quad (12)$$

$$H(\mathbf{x}; r) = \int_{\partial B(\mathbf{x}; r)} \psi(\mathbf{y}, t) d\mathbf{y} \quad (13)$$

For fixed  $\mathbf{x}$ , we regard  $U$  as a function of  $r$  and  $t$ . We claim

**Lemma 1** *For fixed  $\mathbf{x}$ ,  $U(\mathbf{x}; r, t)$  is a solution of the initial value problem:*

$$U_{tt} - U_{rr} - \frac{n-1}{r} U_r = 0 \text{ for } r > 0, t > 0 \text{ and } U(\mathbf{x}; r, 0) = G(\mathbf{x}, r), \quad U_t(\mathbf{x}; r, 0) = H(\mathbf{x}, r) \quad (14)$$

PROOF. For convenience, we depart from our usual convention and denote ‘surface area’ element on the  $n$ -dimensional ball as  $dS$ . Symbol  $dS_{\mathbf{y}}$  will denote surface area element in the variable  $\mathbf{y}$ . We note that

$$U(\mathbf{x}; r, t) = \int_{\partial B(\mathbf{x}, r)} u(\mathbf{y}, t) dS_{\mathbf{y}} = \int_{\partial B(\mathbf{0}, 1)} u(\mathbf{x} + r\mathbf{z}, t) dS_{\mathbf{z}} \quad (15)$$

Therefore,

$$\begin{aligned} U_r(\mathbf{x}; r, t) &= \int_{\partial B(\mathbf{0}, 1)} \mathbf{z} \cdot \nabla u(\mathbf{x} + r\mathbf{z}, t) dS_{\mathbf{z}} = \int_{\partial B(\mathbf{x}, r)} \frac{\partial u}{\partial n} dS_{\mathbf{y}} = \frac{1}{A_r} \int_{\partial B(\mathbf{x}, r)} \frac{\partial u}{\partial n} dS_{\mathbf{y}} \\ &= \frac{1}{A_r} \int_{B(\mathbf{x}, r)} \Delta u(\mathbf{y}, t) d\mathbf{y} = \frac{1}{n\alpha(n)r^{n-1}} \int_{B(\mathbf{x}, r)} \Delta u(\mathbf{y}, t) d\mathbf{y}, \quad (16) \end{aligned}$$

Thus, using (10), it follows that

$$U_r(\mathbf{x}; r, t) = \frac{1}{n\alpha(n)r^{n-1}} \int_{B(\mathbf{x}, r)} u_{tt} d\mathbf{y} \quad (17)$$

and therefore,

$$\frac{\partial}{\partial r} \{r^{n-1}U_r(\mathbf{x}; r, t)\} = \frac{1}{n\alpha(n)} \int_{\partial B(\mathbf{x}; r)} u_{tt}(\mathbf{y}, t) dS_{\mathbf{y}} = r^{n-1}U_{tt} \quad (18)$$

This gives the PDE for  $U$  given in the Lemma. Further, it is clear from definition of  $G$  and  $H$  that  $U$  satisfies the given initial conditions.  $\square$

**Theorem 2** (*Kirchoff Formula for  $n = 3$* )

*The solution to the initial value problem for the three-dimensional dimensional wave equation in free-space:*

$$u_{tt} - \Delta u = 0 \quad \text{for } \mathbf{x} \in \mathbb{R}^3 \text{ for } t > 0 \quad \text{with } u(\mathbf{x}, 0) = \phi(\mathbf{x}) \quad , \quad u_t(\mathbf{x}, 0) = \psi(\mathbf{x}) \quad (19)$$

is given by

$$u(\mathbf{x}, t) = \int_{\partial B(\mathbf{x}, t)} \{t\psi(\mathbf{y}) + \phi(\mathbf{y}) + (\mathbf{y} - \mathbf{x}) \cdot \nabla\phi(\mathbf{y})\} dS_{\mathbf{y}} \quad (20)$$

PROOF.

If note that if we introduce transformation

$$\tilde{U}(\mathbf{x}; r, t) = rU(\mathbf{x}; r, t), \quad \tilde{G} = rG, \quad \tilde{H} = rH$$

Then, simple calculation shows

$$\tilde{U}_{tt} - \tilde{U}_{rr} = 0 \quad \text{for } r > 0 \quad , \quad t > 0, \quad \text{with } \tilde{U}(r, 0) = \tilde{G}(r), \quad \tilde{U}_t(r, 0) = \tilde{H}(r) \quad , \quad \tilde{U}(0, t) = 0$$

This is the Wave equation on a half-line with a homogeneous *Dirichlet* condition. As discussed in Week 4 lectures (see equation (32) on page 4, with  $c = 1$ ) for  $0 \leq r \leq t$ , we obtain

$$\tilde{U}(\mathbf{x}; r, t) = \frac{1}{2} [\tilde{G}(r+t) - \tilde{G}(t-r)] + \frac{1}{2} \int_{t-r}^{r+t} \tilde{H}(y) dy$$

Since  $u(\mathbf{x}, t) = \lim_{r \rightarrow 0^+} \frac{\tilde{U}(\mathbf{x}; r, t)}{r}$ ,

$$\begin{aligned} u(\mathbf{x}, t) &= \lim_{r \rightarrow 0^+} \left\{ \frac{1}{2r} [\tilde{G}(r+t) - \tilde{G}(t-r)] + \frac{1}{2r} \int_{t-r}^{r+t} \tilde{H}(y) dy \right\} \\ &= \tilde{G}'(t) + \tilde{H}(t) = \partial_t \left( t \int_{\partial B(\mathbf{x}, t)} \phi dS \right) + t \int_{\partial B(\mathbf{x}, t)} \psi dS = \partial_t \left( t \int_{\partial B(\mathbf{0}, 1)} \phi(\mathbf{x} + t\mathbf{z}) dS_{\mathbf{z}} \right) + t \int_{\partial B(\mathbf{0}, 1)} \psi dS_{\mathbf{z}} \\ &= \int_{\partial B(\mathbf{0}, 1)} \{t\psi(\mathbf{x} + t\mathbf{z}) + \phi(\mathbf{x} + t\mathbf{z}) + t\mathbf{z} \cdot \nabla\phi(\mathbf{x} + t\mathbf{z})\} dS_{\mathbf{z}} = \int_{\partial B(\mathbf{x}, t)} \{t\psi(\mathbf{y}) + \phi(\mathbf{y}) + \nabla\phi(\mathbf{y}) \cdot (\mathbf{y} - \mathbf{x})\} dS_{\mathbf{y}} \end{aligned}$$

$\square$

**Theorem 3** (Poisson Formula for  $n = 2$ )

The solution to the initial value problem for the two-dimensional dimensional wave equation in free-space:

$$u_{tt} - \Delta u = 0 \text{ for } \mathbf{x} \in \mathbb{R}^2 \text{ for } t > 0 \text{ with } u(\mathbf{x}, 0) = \phi(\mathbf{x}) \text{ , } u_t(\mathbf{x}, 0) = \psi(\mathbf{x}) \quad (21)$$

is given by

$$u(\mathbf{x}, t) = \frac{1}{2} \int_{B(\mathbf{x}, t)} \frac{\{t\psi(\mathbf{y}) + t^2\psi(\mathbf{y} + t(\mathbf{y} - \mathbf{x}) \cdot \nabla\phi(\mathbf{y}))\}}{(t^2 - |\mathbf{y} - \mathbf{x}|^2)^{1/2}} d\mathbf{y} \quad (22)$$

PROOF. We imbed the 2-D problem as part of 3-D problem. With  $\bar{\mathbf{x}} = (x_1, x_2, x_3)$ ,  $\mathbf{x} = (x_1, x_2)$ ,  $\bar{u}(\bar{\mathbf{x}}, t) = u(\mathbf{x}, t)$ . Then  $\bar{u}$  satisfies the 3-D wave equation with initial condition

$$\bar{\phi}(\bar{\mathbf{x}}) = \phi(\mathbf{x}) \text{ and } \bar{\psi}(\bar{\mathbf{x}}) = \psi(\mathbf{x})$$

Then, we have from the 3-D calculation,

$$\bar{u}(\bar{\mathbf{x}}, t) = \partial_t \left( t \int_{\partial\bar{B}(\bar{\mathbf{x}}, t)} \bar{\phi} d\bar{S} \right) + t \int_{\partial\bar{B}(\bar{\mathbf{x}}, t)} \bar{\psi} d\bar{S}$$

where  $\bar{B}(\bar{\mathbf{x}}, t)$  denotes the ball in  $\mathbb{R}^3$  with center  $\bar{\mathbf{x}}$  of radius  $t > 0$ , and  $d\bar{S}$  denotes the two dimensional surface measure. Now we observe that

$$\int_{\partial\bar{B}(\bar{\mathbf{x}}, t)} \bar{g} d\bar{S} = \frac{1}{4\pi t^2} \int_{\partial\bar{B}(\bar{\mathbf{x}}, t)} \bar{g} d\bar{S} = \frac{2}{4\pi t^2} \int_{B(\mathbf{x}, t)} g(\mathbf{y})(1 + (\nabla\gamma)^2)^{1/2} d\mathbf{y}$$

where  $\gamma(\mathbf{y}) = \sqrt{t^2 - |\mathbf{y} - \mathbf{x}|^2}$  for  $\mathbf{y} \in B(\mathbf{x}, t)$ . The factor 2 enters since  $\partial\bar{B}(\bar{\mathbf{x}}, t)$  consists of two hemispheres. Computation shows that  $[1 + (\nabla\gamma)^2]^{1/2} = t[t^2 - |\mathbf{y} - \mathbf{x}|^2]^{-1/2}$ . Therefore,

$$\int_{\partial\bar{B}(\bar{\mathbf{x}}, t)} \bar{g} d\bar{S} = \frac{t}{2} \int_{B(\mathbf{x}, t)} \frac{g(\mathbf{y})}{\sqrt{t^2 - |\mathbf{y} - \mathbf{x}|^2}} d\mathbf{y} = \int_{B(\mathbf{0}, 1)} \frac{g(\mathbf{x} + t\mathbf{z})}{\sqrt{1 - |\mathbf{z}|^2}} d\mathbf{z}$$

The rest of the theorem is straight-forward computation.  $\square$

## 2.2 Source solution for Wave Equation

We consider source solution  $S(\mathbf{x}, t)$  that satisfies:

$$S_{tt} = c^2 \Delta S \text{ for } \mathbf{x} \in \mathbb{R}^n \text{ , } t \in \mathbb{R} \text{ with } S(\mathbf{x}, 0) = 0 \text{ ; } S_t(\mathbf{x}, 0) = \delta(\mathbf{x}) \quad (23)$$

This is referred to as the *Riemann* problem. To find formula for  $S$ , let  $\psi(\mathbf{x})$  be any test function and we define

$$u(\mathbf{x}, t) = \int_{\mathbb{R}^n} S(\mathbf{x} - \mathbf{y}, t) \psi(\mathbf{y}) d\mathbf{y} \quad (24)$$

Then, assuming integration with respect to  $\mathbf{x}$  and  $t$  commutes with the integration with respect to  $\mathbf{y}$ , it follows that  $u$  satisfies wave equation as well, and satisfies initial conditions

$$u(\mathbf{x}, 0) = 0, \text{ and } u_t(\mathbf{x}, 0) = \psi(\mathbf{x}) \quad (25)$$

From D'Alembert formula, the solution to this for  $n = 1$  is given by

$$u(x, t) = \int_{-\infty}^{\infty} S(x - y, t) \psi(y) dy = \frac{1}{2c} \int_{x-ct}^{x+ct} \psi(y) dy$$

Therefore,  $S(x - y, t) = \frac{1}{2c}$  for  $y - x \in (-ct, ct)$  and 0 otherwise. Therefore,

$$S(x, t) = \frac{1}{2c} \text{ for } |x| < ct \text{ and } 0 \text{ for } |x| > ct \text{ for } t > 0 \quad (26)$$

Similar formula can be found for  $t < 0$ . Notice that if we replace  $t$  by  $-t$  in the initial value problem, it only reverses the sign of  $\psi$ . Using Heaviside function  $H^1$ , we obtain

$$S(x, t) = \frac{1}{2c} H(c^2 t^2 - x^2) \text{sgn}(t) \quad (27)$$

For 1-D, the *Riemann* function is actually a function in the usual sense. This is not the case in higher dimension, where it is a distribution.

Note from *Kirchoff*-formula that solution for  $t > 0$  for  $n = 3$  is given by

$$\frac{1}{4\pi c^2 t} \int_{\partial B(\mathbf{x}, t)} \psi(\mathbf{y}) dS_{\mathbf{y}} = u(\mathbf{x}, t) = \int_{\mathbb{R}^3} S(\mathbf{x} - \mathbf{y}, t) \psi(\mathbf{y}) d\mathbf{y} = \int_0^\infty dr \int_{\partial B(\mathbf{x}, r)} S(\mathbf{y} - \mathbf{x}, t) \psi(\mathbf{y}) dS_{\mathbf{y}} \quad (28)$$

Therefore for  $t > 0$ ,  $S(\mathbf{x}, t) = \frac{1}{4\pi c^2 t} \delta(|\mathbf{x}| - ct)$ . We can similarly analyze the solution for  $t < 0$ , noticing that that replacing  $t$  by  $-t$  in the problem posed for  $S$  has the effect of switching its sign. Therefore, for  $t < 0$ , we must have  $S(\mathbf{x}, t) = -\frac{1}{4\pi c^2 (-t)} \delta(|\mathbf{x}| + ct)$ . A uniform expression is given by

$$S(\mathbf{x}, t) = \frac{1}{2\pi c} \delta(|\mathbf{x}|^2 - c^2 t^2) \text{sgn}(t)$$

In 2-D similar calculation using Poisson formula shows

$$S(\mathbf{x}, t) = \frac{1}{2\pi c} (c^2 t^2 - |\mathbf{x}|^2)^{-1/2} \text{ for } |\mathbf{x}| < ct \text{ and } = 0 \text{ otherwise}$$

### 3 Eigen Function Expansion for Green's Function

#### 3.1 Heat Equation

Consider Source solution to heat equation bounded domain  $\Omega \subset \mathbb{R}^n$  with homogeneous Dirichlet Boundary conditions:

$$S_t = \kappa \Delta S \text{ for } \mathbf{x} \in \Omega \text{ and } S = 0 \text{ on } \partial\Omega, \text{ with } S(\mathbf{x}, 0) = \delta(\mathbf{x} - \mathbf{x}_0) \quad (29)$$

In terms of the  $S(\mathbf{x}, \mathbf{x}_0, t)$  the solution to the initial value problem

$$u_t = \kappa \Delta u \text{ for } \mathbf{x} \in \Omega \text{ and } u = 0 \text{ on } \partial\Omega, \text{ with } u(\mathbf{x}, 0) = \phi(\mathbf{x}) \quad (30)$$

---

<sup>1</sup>Recall  $H(x) = 1$  for  $x > 0$  and  $H(x) = 0$  for  $x < 0$

is given by

$$u(\mathbf{x}, t) = \int_{\Omega} S(\mathbf{x}, \mathbf{y}, t) \phi(\mathbf{y}) d\mathbf{y} \quad (31)$$

On the otherhand, if we denote the orthonormalized eigenfunctions  $\{X_n\}_{n=1}^{\infty}$  and corresponding eigenvalues  $\{\lambda_n\}_{n=1}^{\infty}$  of the operator  $-\Delta$  with homogenous boundary conditions on  $\partial\Omega$ , we know that solution to heat equation has the form

$$u(\mathbf{x}, t) = \sum_{n=1}^{\infty} \exp[-\lambda_n \kappa t] X_n(\mathbf{x}) \quad (32)$$

where

$$c_n = (\phi, X_n) = \int_{\Omega} \phi(\mathbf{x}) X_n(\mathbf{x}) d\mathbf{x} \quad (33)$$

Then,

$$u(\mathbf{x}, t) = \sum_{n=1}^{\infty} \left( \int_{\Omega} \phi(\mathbf{y}) X_n(\mathbf{y}) d\mathbf{y} \right) \exp[-\lambda_n \kappa t] X_n(\mathbf{x}) = \left( \int_{\Omega} \phi(\mathbf{y}) \left\{ \sum_{n=1}^{\infty} X_n(\mathbf{y}) X_n(\mathbf{x}) e^{-\lambda_n \kappa t} \right\} d\mathbf{y} \right) \quad (34)$$

Therefore, it follows that under the assumption that the summation converges absolutely and uniformly,

$$S(\mathbf{x}, \mathbf{y}, t) = \sum_{n=1}^{\infty} X_n(\mathbf{y}) X_n(\mathbf{x}) e^{-\lambda_n \kappa t} \quad (35)$$

Note, that this implies that in the sense of distribution, we must have

$$\delta(\mathbf{x} - \mathbf{y}) = \sum_{n=1}^{\infty} X_n(\mathbf{y}) X_n(\mathbf{x})$$

This is true for any complete ortho-normal basis.

## 4 Fourier Transform

**Notation:** For multi-index  $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_n)$ , with each  $\alpha_j \in \mathbb{N}$ , it is convenient to introduce operator

$$D^{\alpha} \equiv \partial_{x_1}^{\alpha_1} \partial_{x_2}^{\alpha_2} \dots \partial_{x_n}^{\alpha_n}$$

The order of this operator is denoted by  $|\alpha| \equiv \alpha_1 + \alpha_2 + \dots + \alpha_n$ .

**Definition 4**  $\mathcal{S}(\mathbb{R}^n)$  be the space of all functions  $\phi$  on  $\mathbb{R}^n$  which are of the class  $\mathbf{C}^{\infty}$  and such that for any integer  $j \geq 0$ ,  $|\mathbf{x}|^j |D^{\alpha} \phi| < \infty$ , for  $|\alpha| = j$ . This is referred to usually as the Schwartz class of functions.

**Definition 5** A tempered distribution in  $\mathbb{R}^n$  is a continuous linear functional on the class of  $\phi \in \mathcal{S}(\mathbb{R}^n)$ .

**Remark 1** Note that every tempered-distribution is a distribution, but the converse is not true.

The Fourier transform of a continuous absolutely integrable function  $f$  on  $\mathbb{R}^n$  is defined by

$$\hat{f}(\mathbf{k}) = \mathcal{F}[f](\mathbf{k}) = (2\pi)^{-n/2} \int_{\mathbb{R}^n} \exp[-i\mathbf{k} \cdot \mathbf{x}] f(\mathbf{x}) d\mathbf{x}$$

In particular, this defines Fourier-Transform for every  $f \in \mathcal{S}(\mathbb{R}^n)$ .

**Theorem 6** *If  $f \in \mathcal{S}(\mathbb{R}^n)$ , then  $\hat{f} \in \mathcal{S}(\mathbb{R}^n)$ . Moreover, the mapping is continuous from  $\mathcal{S}(\mathbb{R}^n)$  to itself.*

PROOF. We leave the proof to the reader.  $\square$

**Theorem 7** *Let  $g \in \mathcal{S}(\mathbb{R}^n)$ . Then there is a unique  $f \in \mathcal{S}(\mathbb{R}^n)$  such that  $g = \mathcal{F}[f]$ . Furthermore, the inverse Fourier transform of  $g$  is given by*

$$f(\mathbf{x}) = (2\pi)^{-n/2} \int_{\mathbb{R}^n} e^{i\mathbf{k} \cdot \mathbf{x}} g(\mathbf{k}) d\mathbf{k} \quad (36)$$

PROOF. Let  $Q_M = [-M, M]^n$ , and let  $f$  be given by the above formula. Then, we find

$$\begin{aligned} \tilde{f}(\mathbf{k}) &= (2\pi)^{-n/2} \int_{\mathbb{R}^n} e^{-i\mathbf{k} \cdot \mathbf{x}} f(\mathbf{x}) d\mathbf{x} = (2\pi)^{-n/2} \int_{\mathbb{R}^n} e^{-i\mathbf{k} \cdot \mathbf{x}} \int_{\mathbb{R}^n} e^{i\boldsymbol{\eta} \cdot \mathbf{x}} g(\boldsymbol{\eta}) d\boldsymbol{\eta} d\mathbf{x} \\ &= \lim_{M \rightarrow \infty} \int_{\mathbb{R}^m} \int_{Q_M} e^{i(\boldsymbol{\eta} - \mathbf{k}) \cdot \mathbf{x}} g(\boldsymbol{\eta}) d\mathbf{x} d\boldsymbol{\eta} = \pi^{-n} \lim_{M \rightarrow \infty} \int_{\mathbb{R}^m} \prod_{i=1}^n \frac{\sin M(\eta_i - k_i)}{\eta_i - k_i} g(\boldsymbol{\eta}) d\boldsymbol{\eta} \end{aligned}$$

However, it is easily seen that as  $M \rightarrow \infty$ ,  $\frac{\sin M(\eta_i - k_i)}{\eta_i - k_i} \rightarrow \pi \delta(\eta_i - k_i)$  in the sense of distribution. Therefore, it follows that

$$\hat{f}(\mathbf{k}) = g(\mathbf{k})$$

An analogous calculation shows that if  $g = \hat{h}$  for some  $h \in \mathcal{S}(\mathbb{R}^m)$ , then  $h = f$  as given by equation (36).  $\square$

**Theorem 8** *Let  $f, g \in \mathcal{S}(\mathbb{R}^n)$ , then  $(\hat{f}, \hat{g}) = (f, g)$ .*

PROOF. We have

$$\begin{aligned} (f, g) &= \int_{\mathbb{R}^n} f(\mathbf{x}) \overline{g(\mathbf{x})} d\mathbf{x} = (2\pi)^{-n/2} \int_{\mathbb{R}^n} \overline{g(\mathbf{x})} \int_{\mathbb{R}^n} \hat{f}(\mathbf{k}) e^{i\mathbf{k} \cdot \mathbf{x}} d\mathbf{k} d\mathbf{x} \\ &= (2\pi)^{-n/2} \int_{\mathbb{R}^n} \hat{f}(\mathbf{k}) \int_{\mathbb{R}^n} \overline{g(\mathbf{x})} e^{-i\mathbf{k} \cdot \mathbf{x}} d\mathbf{x} d\mathbf{k} = \int_{\mathbb{R}^n} \overline{\hat{g}(\mathbf{k})} \hat{f}(\mathbf{k}) d\mathbf{k} = (\hat{f}, \hat{g}) \end{aligned}$$

$\square$

We now seek to give meaning to Fourier-Transform of tempered distribution.

**Definition 9** *Let  $f \in \mathcal{S}'(\mathbb{R}^n)$ . Then the Fourier transform of  $f$  is defined by the functional*

$$(\mathcal{F}[f], \phi) = (f, \mathcal{F}^{-1}[\phi]) \quad \text{for } \phi \in \mathcal{S}(\mathbb{R}^n)$$

**Remark 2** It is not difficult to see that  $\mathcal{F}$  is a continuous mapping from  $\mathcal{S}'(\mathbb{R}^n)$  onto itself. The formulas for Fourier transform and its inverse still hold for tempered distribution.

**Example** We want Fourier-transform of  $\delta$  distribution. From definition

$$(\mathcal{F}[\delta], \phi) = (\delta, \mathcal{F}^{-1}[\phi]) = \mathcal{F}^{-1}[\phi](\mathbf{0}) = (2\pi)^{-n/2} \int_{\mathbb{R}^n} \phi(\mathbf{x}) d\mathbf{x}$$

Therefore  $\mathcal{F}[\delta] = (2\pi)^{-n/2}$ , a constant.

**Example** The above relation is symmetric. since the Fourier-transform of 1 equal to  $(2\pi)^{n/2}\delta$  since

$$(\mathcal{F}[1], \phi) = (1, \mathcal{F}^{-1}[\phi]) = \int_{\mathbb{R}^n} \mathcal{F}^{-1}[\phi](\mathbf{x}) d\mathbf{x} = (2\pi)^{n/2} \mathcal{F}\mathcal{F}^{-1}[\phi](\mathbf{0}) = (2\pi)^{n/2} \phi(\mathbf{0})$$

Therefore,  $\mathcal{F}[1] = (2\pi)^{n/2}\delta$

**Example:** Let  $\delta(|\mathbf{x}| - a)$  represent a uniform mass distribution on a sphere of radius  $a$ , i.e.

$$(\delta(|\mathbf{x}| - a), \phi) = \int_{|\mathbf{x}|=a} \phi(\mathbf{x}) dS$$

Then,

$$\mathcal{F}[\delta(|\mathbf{x}| - a)](\mathbf{k}) = (2\pi)^{-n/2} \int_{|\mathbf{x}|=a} e^{-i\mathbf{k}\cdot\mathbf{x}} dS$$

For  $n = 3$ , using spherical polar coordinates, we get

$$\mathcal{F}[\delta(|\mathbf{x}| - a)](\mathbf{k}) = (2\pi)^{-n/2} \int_0^\pi \int_0^{2\pi} e^{-ia\rho \cos\theta} \sin\theta d\phi d\theta = \sqrt{\frac{2}{\pi}} a \frac{\sin a|\mathbf{k}|}{|\mathbf{k}|}$$